

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/335732294>

Controlled Environment Food Production for Urban Agriculture

Article in HortScience: a publication of the American Society for Horticultural Science · September 2019

DOI: 10.21273/HORTSCI14073-19

CITATIONS

6

READS

1,228

12 authors, including:



Christopher J. Currey
Iowa State University

60 PUBLICATIONS 359 CITATIONS

[SEE PROFILE](#)



Hye-Ji Kim
Purdue University

60 PUBLICATIONS 429 CITATIONS

[SEE PROFILE](#)



Nadia Sabeh
Dr. Greenhouse

20 PUBLICATIONS 168 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Plant Factory [View project](#)



Climate Science Communication to Rural Audiences [View project](#)

Controlled Environment Food Production for Urban Agriculture

Celina Gómez²

Environmental Horticulture Department, University of Florida, 1549 Fifield Hall, Gainesville, FL 32611

Christopher J. Currey

Department of Horticulture, Iowa State University, 008 Horticulture Hall, Ames, IA 50011

Ryan W. Dickson

Department of Horticulture, University of Arkansas, PTSC 311, Fayetteville, AR 72701

Hye-Ji Kim

Department of Horticulture and Landscape Architecture, Purdue University, 625 Agriculture Mall Drive, West Lafayette, IN 47907

Ricardo Hernández

Department of Horticultural Sciences, North Carolina State University, 2721 Founders Drive, Suite 170, Raleigh, NC 27695

Nadia C. Sabeh

Dr. Greenhouse, Inc., 930 Alhambra Boulevard, Suite 260, Sacramento, CA 95816

Rosa E. Raudales

Department of Plant Science and Landscape Architecture, University of Connecticut, 1376 Storrs Road, Storrs CT 06269

Robin G. Brumfield

Department of Agriculture, Food and Resource Economics, Rutgers University, 55 Dudley Road, New Brunswick, NJ 08901

Angela Laury-Shaw

Department of Food Science and Human Nutrition, Iowa State University, 2577 Food Sciences Building, 536 Farm House Lane, Ames, IA 50011

Adam K. Wilke¹

Iowa State University Extension and Outreach, 418 East Hall, Ames, IA 50011

Roberto G. Lopez

Department of Horticulture, Michigan State University, 1066 Bogue Street, East Lansing, MI 48824

Stephanie E. Burnett

School of Food and Agriculture, The University of Maine, 5722 Deering Hall, Orono, ME 04469

Additional index words. economics, food safety, food security, greenhouse, plant factory, review, sole-source lighting, and supplemental lighting

Abstract. The recent increased market demand for locally grown produce is generating interest in the application of techniques developed for controlled environment agriculture (CEA) to urban agriculture (UA). Controlled environments have great potential to revolutionize urban food systems, as they offer unique opportunities for year-round production, optimizing resource-use efficiency, and for helping to overcome significant challenges associated with the high costs of production in urban settings. For urban growers to benefit from CEA, results from studies evaluating the application of controlled environments for commercial food production should be considered. This review includes a discussion of current and potential applications of CEA for UA, references discussing appropriate methods for selecting and controlling the physical plant production environment, resource management strategies, considerations to improve economic viability, opportunities to address food safety concerns, and the potential social benefits from applying CEA techniques to UA. Author's viewpoints about the future of CEA for urban food production are presented at the end of this review.

The term controlled-environment agriculture (CEA) was first introduced in the 1960s and refers to an intensive approach for controlling plant growth and development by capitalizing on advanced horticultural techniques and innovations in technology (Hodges et al., 1968). Controlled environments (CEs) provide advantages to predict plant responses to their environment and increase production efficiency, optimize plant yield, and improve product quality. They play a key role in the commercial production of ornamental plants and vegetable crops and in the production of young plant material from seed, cuttings, or tissue-culture (Jensen, 2002; Kozai and Niu, 2016a). The recent increase in market demand for locally grown produce is generating interest in applying CEA practices to urban agriculture (UA), including small- (e.g., in-home production or indoor gardens), medium- (e.g., community gardens), or large-scale commercial operations [e.g., rooftop greenhouses or warehouse-based indoor “plant factories” (PFs), sometimes referred to as vertical, warehouse, or container farms] (Eaves and Eaves, 2018; Jansen et al., 2016).

Before land and labor shortages prompted by the Industrial Revolution forced food production to move away from cities, agriculture was central to urban environments and their planning (Vitiello and Brinkley, 2014). Although some efforts were made to promote UA by documenting health, educational, and social benefits, urban food production was viewed mostly as a strategy to reduce pressure on the public food supply during hardships such as war (Hynes and Howe, 2004; Lawson, 2005). However, recent shifts in consumption patterns are allowing UA to make a comeback through its concerted efforts to address sustainability issues in our food system (e.g., reduced dependence on fossil fuels, increased food security) and promote social and environmental cohesion (Peterson et al., 2015). CEs have tremendous potential for commercial urban food production. The high plant density and year-round production attainable with CEA optimize space use and can help overcome challenges associated with costs and availability of land in urban settings. Moreover, soilless culture systems allow plants to grow in nonconventional spaces, as opposed to fertile soil, while maximizing use of available resources (e.g., water and nutrients). For a comprehensive review of the benefits from CEA in UA, see Kozai and Niu (2016b) and Takagaki et al. (2016).

For UA producers to benefit from CEA, results from studies evaluating the applicability of CEs for commercial food production should be considered. Although profitability will typically depend on the local demand and supply of food, location, population density, facility design, and crops produced, preliminary research suggests that economic sustainability for commercial CEA requires careful consideration of capital investment and operating costs, production volume, product quality and consistency, and local market trends (Al-Kodmany, 2018; Kozai et al., 2016). A review of key studies, followed by

a discussion of current and potential applications regarding opportunities and limitations of commercial CEA for urban food production follows.

Physical Environment

Production environment. A key benefit of CEA is the ability to modify production environments to maximize plant quality and yield, extend growing seasons, and enable crop production in unfavorable climatic conditions (e.g., wind, rain, extreme temperature, and limited light). Greenhouses and PFs are the most common types of CEs used in UA. Following the construction of an efficient transportation system after World War II, most commercial greenhouses in the United States settled in rural locations with favorable climates (e.g., high-light, moderate temperatures) or near large markets to improve economic viability (Nelson, 2012). Therefore, it is not uncommon for greenhouse-grown food crops to travel long distances before reaching consumers (Weber and Matthews, 2008). However, increasing interest in local food systems is bringing attention to the impact that greenhouse gas emissions associated with food production against long-distance distribution channels (i.e., “food miles”) have on our food chain. Although some argue that reducing “food miles” is necessary to improve the sustainability of our food chain (Morgan, 2009; Paxton, 1994), others call attention to the fact that several key transportation factors (e.g., vehicle efficiency, infrastructure, alternative fuels) affect the sustainability of many CEs typically used in UA (Coley et al., 2009; Mundler and Rumpus, 2012). However, the potential for reducing environmental impacts from UA needs to be assessed from a broad perspective, including evaluating the life cycle assessment (LCA) of a product and incorporating socioeconomic factors (Edwards-Jones et al., 2008).

Greenhouses built on vacant rooftops of city buildings have become popular in recent years as they capitalize on sunlight to produce plant products in close proximity to consumers. In addition, because rooftops represent close to one fourth of residential and nonresidential urban areas, they have significant potential to contribute to the expansion of UA (Getter and Rowe, 2006). Nonetheless, important economic aspects of producing in rooftop greenhouses should be considered, as they may require special construction materials, components, and struc-

tural support to satisfy building codes and withstand strong wind loads and substantial sun exposure (Meier et al., 2013). Numerous studies assess the impact of building rooftop greenhouses in urban settings from an economic (Benis et al., 2018; Sanyé-Mengual et al., 2015), environmental (Jones and Gilbert, 2018; Nadal et al., 2017; Sanjuan-Delmás et al., 2018; Sanyé-Mengual et al., 2018), social (Davis, 2011), and educational (Nadal et al., 2017; Sanyé-Mengual et al., 2014) perspective. In addition, a number of commercial operations have demonstrated the economic viability of rooftop greenhouses for urban food production (e.g., Gotham Green Farms, Lufa Farms). Furthermore, rooftop greenhouse designs can include photovoltaic systems and rainwater harvesting strategies, increasing their potential to expand sustainable UA production practices.

In contrast, the economic sustainability of indoor PFs has demonstrated to be more challenging, in part due to their dependence on electricity to run all systems (Banerjee and Adenauer, 2014). Indoor PFs are widely used in Asia and are gaining popularity in the United States and European countries for producing high-value crops such as leafy greens (Kozai et al., 2016). However, most countries successfully adopting PFs for fresh food production have land and/or environmental limitations, which, coupled with food-safety concerns, justify the high production costs. However, preliminary research suggests that some U.S. consumers are not willing to pay higher prices for indoor-grown produce (Coyle and Ellison, 2017; Short et al., 2018).

Production systems. Production systems for CEA typically consist of soilless culture, with or without the use of an organic or inorganic substrate and with active application of water and fertilizer, typically provided with a dilute nutrient solution. Hydroponic production is a type of soilless culture in which plant roots are suspended in either a static, continuously aerated nutrient solution, or in a continuous flow or mist (Jones, 2014). The terms “soilless culture” and “hydroponics” often are used interchangeably, especially when using substrates that are chemically inert and provide little nutrient supply or retention.

Crop production using soilless culture or hydroponics has several potential benefits compared with traditional field production, including isolation from soil or water-borne issues (e.g., nematodes, salinity, or heavy metals) and an improved ability to control water and nutrient uptake. For comprehensive reviews about the benefits of using soilless culture in CEA, see Raviv et al. (2019) and Jones (2014). Crops produced using soilless culture often are grown in containers, troughs, or bags with a limited volume, allowing for efficient root-zone management. Argo and Fisher (2002), Bunt (1988), and Raviv et al. (2019) list common sources of nutrients in soilless culture, which include the raw irrigation water, fertilizers either dissolved in the irrigation water or incorporated into a sub-

strate, substrate components, and amendments used to adjust substrate pH. Raviv et al. (2019) also discuss common irrigation and fertilization equipment used for container production systems, including overhead, drip, and sub-irrigation systems. Nelson (2012) provides a detailed review of management strategies across soilless culture systems, including strategies on system design, substrates, water and fertilizer, production environment, and marketing and business planning.

Substrate selection is a critical aspect of soilless culture. Primary substrate components (i.e., >40% of the substrate volume) often consist of organic materials with low bulk density and high water holding capacity, such as peatmoss and coconut coir fiber (Argo and Fisher, 2002). Conversely, secondary components (i.e., <40% substrate volume) often include materials such as expanded minerals (e.g., perlite and vermiculite), clays, sand, and composts that increase drainage and cation exchange capacity to increase aeration and nutrient retention. Several authors have reviewed the most common organic and inorganic components of substrates, manufacturing, and cultural plant production advantages and disadvantages to using different materials (Bunt, 1988; Burnett et al., 2016; Raviv et al., 2019).

Hydroponic systems commonly used in UA differ in their design and suitability for certain crops and production scenarios. The most common hydroponic systems include nutrient-film technique (NFT), deep-water culture (DWC; also known as deep-flow technique, raft, raceway, or floating hydroponics, among others), and aggregate culture. NFT consist of crops grown in sloped troughs where a thin film of nutrient solution flows (either continuously or intermittently) over the roots, whereas DWC systems consist of crops grown with their roots continuously submerged in a nutrient solution. Aggregate culture consists of growing crops in bagged substrates (e.g., rockwool or coconut coir slabs) or containers (e.g., Dutch/Bato buckets) with the nutrient solution applied using drip emitters. Typically, NFT and DWC systems are used for short-term, non-fruiting crops such as leafy greens and herbs, whereas long-term fruiting crops such as tomato (*Lycopersicon esculentum*), cucumber (*Cucumis sativus*), sweet pepper (*Capsicum annuum*), and strawberry (*Fragaria ×ananassa*) are usually grown in aggregate culture. Several authors have reviewed specific plant species, water and nutrient management, structural design, and economic considerations for producing crops in NFT and DWC (Blok et al., 2017; Walters and Currey, 2015), and aggregate culture (Raviv et al., 2019; Sonneveld and Voogt, 2009).

Soilless culture and hydroponic systems used in CEs also provide opportunities for producing medicinal and pharmaceutical crops with high quality, purity, consistency, bioactivity, and biomass (Hayden, 2006; Lopez and Runkle, 2017; Papadopoulos et al., 2001; Potter, 2014). Hayden (2006)

Received for publication 22 Mar. 2019. Accepted for publication 22 May 2019.

We gratefully acknowledge support from the US Department of Agriculture National Institute of Food and Agriculture, Multistate Research Project NE1835: Resource Optimization in Controlled Environment Agriculture and thank Dr. A.J. Both for his valuable review.

¹Current address: University of Minnesota Water Resources Center, 183 McNeal Hall, 1985 Buford Avenue, St. Paul, MN 55108.

²Corresponding author: E-mail: comezv@ufl.edu.

and Maggini et al. (2014) discuss the advantages of using hydroponic systems for producing medicinal crops commonly grown in CEs. Moreover, despite being regulated as an illicit crop in many countries, production of cannabis (*Cannabis sativa*) using soilless culture in CEs has increased and gained significant public attention in recent years, particularly in Europe and North America. Cultivation, processing, and regulation of cannabis for medicinal use is reviewed by Potter (2014).

Aquaponic production, integrating hydroponic systems with aquaculture (i.e., fish production), recently has increased in popularity for UA (dos Santos, 2016). Aquaponics involves converting water and organic waste produced by cultivated fish or crustaceans into nutrient solutions used for hydroponic plant production. Potential benefits of aquaponics include the ability to increase resource-use efficiency (e.g., shared startup, operating, and infrastructure costs) and improve sustainability (e.g., reduced water use and waste discharge to the environment) while simultaneously producing fish and food crop commodities (Tyson et al., 2011). Recirculating the nutrient solution between the hydroponic and aquaculture production components, essentially forming a completely closed system, is termed “coupled” aquaponics. Due to the technical challenges with maintaining appropriate chemical and biological properties in a recirculating solution, “decoupled” aquaponics has been proposed as a more efficient alternative to “coupled” aquaponics. In “decoupled” aquaponics, the aquaculture effluent is collected and supplemented with specific nutrients for higher yield of fish and hydroponic food crops (Kloas et al., 2015). However, commercial applications of “decoupled” aquaponics are still under development. Although integrated aquaponic production systems have been used for over 30 years, the commercial application of aquaponics is relatively new to UA. For a comprehensive review on the history of aquaponics see Palm et al. (2018).

For appropriate functioning, aquaponic systems require balancing multiple factors, including solution pH (Wongkiew et al., 2017a; Zou et al., 2016), plant and animal density (Buzby and Lin, 2014; Hussain et al., 2014, 2015), solution flow rates (Hussain et al., 2015; Khater and Ali, 2015; Wongkiew et al., 2017b), economics (Quagraine et al., 2018; Tokunaga et al., 2015), system configuration (Klemencic and Bulc, 2015; Monsees et al., 2017), and food safety considerations (Elumalai et al., 2017; Pantanella et al., 2015). Several authors have reviewed system types, management, and potential profitability of aquaponic production systems (Blidariu and Grozea, 2011; Lewis et al., 1978; Love et al., 2015).

Leaf chlorosis and/or reduced yields are reported for some crops grown in aquaponics, such as tomato (unpublished data) and eggplant (*Solanum melongena*) (Roosta and Mohsenian, 2015). However, supplementing aquaponic solutions with potassium, sulfur,

iron, and manganese may alleviate crop growth and yield losses associated with nutritional deficiencies (Rakocy et al., 2004, 2006; Roosta and Hamidpour, 2011).

Technologies

Electric lighting. The daily light integral (DLI) requirement of food crops commonly grown in CEs typically ranges from 12 to 30 mol·m⁻²·d⁻¹ (Dorais et al., 2017). Greenhouse production located in regions with considerable seasonal variation in solar radiation typically relies on supplemental lighting to increase DLI (Faust and Logan, 2018). Furthermore, solar spectrum can vary depending on location, season, and time of day; however, in general, it is composed of 0.1% ultraviolet-B (280–315 nm), 5% ultraviolet-A (315–400 nm), 19% blue (400–500 nm), 25% green (500–600 nm), 26% red (600–700 nm), 25% far-red (700–800 nm) (extracted from Kotilainen et al., 2018). High-pressure sodium (HPS) lamps are the most common type of electric light source used in commercial greenhouses, whereas metal halide lamps are used occasionally because of their higher blue light output (Menard et al., 2006; Nelson and Bugbee, 2014). However, the use of light-emitting diodes (LEDs) in CEs has substantially increased in recent years (Lopez and Runkle, 2017; Mitchell et al., 2015).

Research comparing LED and HPS lamps as supplemental lighting sources mainly focuses on: 1) evaluating potential reductions in energy consumption; and 2) understanding the effects of light spectra on plant growth, morphology, and quality. Although HPS lamps have a set spectrum that typically is composed of approx. 5% blue, 49% green, 39% red, and 7% far-red (measurements from a 600-W HPS fixture; P.L. Light Systems Inc., Beamsville, ON, Canada), LED fixtures can have different color diodes to achieve customized spectra. However, the most common LED fixtures available for plant lighting contain mainly red and blue diodes (10% to 25% blue diodes), with some fixtures including a small number of far-red and/or white diodes. For a review of different uses, fixture characteristics, and application of LEDs for supplemental lighting in CEA see Lopez and Runkle (2017) and Mitchell et al. (2015).

Although initial efforts to produce high-value food crops in warehouse-based PFs used water-cooled HPS lamps, challenges associated with their economic and thermal management negate the application of these lamps for sole-source lighting in multitiered PFs (Mitchell and Stutte, 2015). Fluorescent lamps have been commonly used for indoor CEA for many years. However, with increasing light efficacy (μmol·J⁻¹), capital cost reductions, and widespread availability, LEDs are becoming the light source of choice for PFs. In fact, introducing commercial LEDs as plant lighting sources was a major contributing factor to the renewed interest in CEA for urban food production. This is

because LED fixtures typically have lower electric power requirements per unit of growing area (kW·m⁻²) and deliver high light intensities with small amounts of radiant heat delivered to crops, theoretically resulting in significant savings for energy-intensive indoor food production. In addition, by maximizing photon capture efficiency using “precision” or “smart” lighting” such as targeted (Poulet et al., 2014), intracanopy (Dueck et al., 2012; Gómez and Mitchell, 2016; Massa et al., 2005), or dynamic (Clausen et al., 2015; Pinho et al., 2012; van Iersel, 2017; van Iersel and Gianino, 2017; van Iersel et al., 2016; Weaver et al., 2019) LED lighting, efficiency of production systems can significantly increase. For a comprehensive review of the adoption of LEDs in UA see Gupta (2017).

Carbon dioxide (CO₂) enrichment. In closed environments with high planting densities (typical for greenhouses and PFs), the CO₂ concentration can rapidly drop below the ambient concentration (approx. 400 μmol·mol⁻¹), requiring supplemental CO₂ injection to avoid limiting photosynthesis and plant growth. Enrichment up to 800 μmol·mol⁻¹ is typically cost-effective to promote photosynthesis and growth of most plants grown in CEs (Both et al., 2017). However, enrichment strategies and benefits depend on, among others: plant species, target CO₂ concentration, environmental control strategies for light, ambient temperature, and humidity, and the cost and source of CO₂ injection (e.g., equipment and electricity) (Both et al., 2017). Indoor CO₂ concentration can be increased by releasing pure gas or by producing CO₂ from fuel combustion. When using the latter approach, carbon monoxide and ethylene production are also important factors to consider with CO₂ enrichment, as they can be toxic to plants at relatively low concentrations and may result from inefficient fuel combustion or by improperly adjusted burners (Nelson, 2012).

Several studies report responses of different food crops to CO₂ enrichment. Generally, studies suggest that doubling ambient CO₂ levels increases lettuce yield by ≈25 to 60% (Chagvardieff et al., 1994; Pérez-López et al., 2015). Conversely, some report small or non-significant differences in lettuce fresh and dry mass grown under elevated CO₂ (>800 ppm) compared with ambient levels (Fu et al., 2015; Mortensen, 1994; Park and Lee, 2001). Yield (kg per plant) of strawberry plants grown in CEs significantly increases by doubling the ambient CO₂ concentration (Enoch et al., 1976; Sun et al., 2012). For cucumber, supplemental nitrate enhances the positive effects of CO₂ enrichment on fruit yield (Dong et al., 2017; Enoch et al., 1976). However, Peet (1986) found no response to cucumber production under an elevated CO₂ concentration. Similarly, although some studies report tomato yield increases with CO₂ enrichment (Calvert and Slack, 1975), others have reported increased individual fruit mass and number of high-rated fruits, but no increase in total yield (Peet and Willits, 1984).

Heuvelink and Kierkels (2015) reported that in general, CO₂ enrichment from 400 to 1000 μmol·mol⁻¹ in CEs can increase total yield by 35 to 50%. For comparison, yield increases with higher DLIs are often linear, as commonly observed in yield differences between greenhouse crops produced during spring-to-summer vs. fall-to-winter seasons (Acock et al., 1971; Cockshull, 1992; Kubota et al., 2016). Also, responses to CO₂ enrichment are species-specific and concentrations above 1000 μmol·mol⁻¹ are not typical for commercial production, since several studies have shown little or no additional benefit at higher concentrations (Calvert and Slack 1975; Enoch et al., 1976; Peet and Willits, 1984). Furthermore, crops with high harvest indices (dry mass of harvestable organs/total plant biomass of plant) such as leafy greens benefit most from CO₂ enrichment, as increases in photosynthesis directly affect the harvestable portion of the plant (i.e., leaves). In contrast, benefits from CO₂ supplementation are less significant with fruiting crops because increased photosynthesis does not directly translate into increased fruit yield.

Humidity control. In CEs, the areal humidity (both absolute and relative) is affected by several factors, such as plant transpiration rate, irrigation strategy, infiltration and ventilation, and active systems used to humidify or dehumidify the air (von Caemmerer and Baker, 2007). Managing humidity is essential for optimal plant growth and development, but it is often the hardest environmental parameter to fully control in CEs. Although relative humidity is the most common measure of humidity, it is not indicative of the plant's relationship to humidity. Vapor pressure deficit (VPD) is the driving force of water loss from a leaf, and therefore, it is a more accurate measure of humidity in CEs. A review of the thermodynamic properties of moist air and their effect on plant growth in CEs can be found in Hanan (1997) and Kozai et al. (2016).

Plant transpiration generally increases linearly with increasing VPD, radiation, and air speed. Therefore, all factors indirectly affect nutrient uptake, leaf temperature, and overall plant growth and development. For leafy greens and culinary herbs grown in CEs, high VPDs are desirable to prevent calcium deficiencies such as tipburn (Frantz et al., 2004; Goto and Takakura, 1992). For CE-grown tomato, VPDs in the range of 0.5 to 1.0 kPa are positively correlated with increases in fruit yield and stomatal conductance, and reductions of blossom-end rot and fungal diseases (Barker, 1990; Guichard et al., 2005; Leonardi et al., 1999; Shamshiri et al., 2018; Zhang et al., 2017). Studies also report that low nighttime VPDs enhance calcium uptake of young and inner strawberry leaves (Bradfield and Guttridge, 1979; Choi et al., 1997), whereas VPDs <0.1 kPa for at least 3 h each night help prevent calcium deficiencies in strawberries produced under environments with high daytime VPD (Kroggel and Kubota, 2017).

Under high-humidity conditions in CEs, water vapor has to be removed from the air to maintain desirable humidity levels. The lowest-cost, lowest-energy method of dehumidification is ventilation, which is the exchange of indoor air with outdoor air, either with fans or vents (Wang et al., 2016). Ventilation works best when outside air is dry and cool. Therefore, in northern climates during winter conditions, ventilation is mainly used for dehumidification. However, if the outside air is too cold, simultaneous heating is needed to maintain the desired air temperature. Alternatively, if the outside air is too hot, cooling will be needed, usually with an air conditioner (common practice for PFs) (Kozai et al., 2016). Growers typically make economic decisions to balance the cost of temperature control with the benefits of lower humidity obtained by ventilation. Kubota et al. (2006) reports that under typical semiarid midday weather, humidity control inside the greenhouse is very efficient. In contrast, humidity control in greenhouses located in continental climates is challenging in the winter months when using ventilation (Kubota et al., 2006). Therefore, the common humidity set point in greenhouses located in continental climates is 80 to 85% relative humidity (Körner and Challa, 2003).

Ventilation is not desirable in PF operations due to reductions in CO₂-use-efficiency from venting out supplemental CO₂ and the potential introduction of pests from the outdoors (Kozai, 2013). For that reason, dehumidification with air conditioning is typically used in PFs, as opposed to ventilation in greenhouses. However, reports using an internal dehumidification system in the greenhouse (dehumidifier or heat pump) indicate reductions in energy consumption compared with the venting-heating technique (de Zwart, 2014; Vallières et al., 2014). Furthermore, several studies have evaluated night-time greenhouse dehumidification using a heat pump to prevent condensation on foliage (Boulard et al., 1989; Campen et al., 2003; Chassériaux and Gaschet, 2011; Migeon et al., 2012), preventing venting the greenhouse during the night to reduce the consumption of energy by conditioning (re-heating) the air. In some instances, night-time heat-pump dehumidification requires up to five times less energy than the common venting-heating technique (Chassériaux et al., 2014). Furthermore, proof-of-concept studies demonstrate the feasibility of using desiccant-based systems to absorb moisture from the greenhouse environment during periods with high humidity (Davies, 2005; Lychnos and Davies, 2012; Mei and Dai, 2008).

Transpiration increases the amount of water in the air and increases the energy content (i.e., enthalpy) of the air. Heating, ventilation, and air conditioning systems are able to cool the air (remove energy) while lowering the humidity (dehumidification). In PFs, it is not uncommon for the energy input from transpiration (i.e., latent heat) to be larger than the energy input from electric

equipment (i.e., sensible heat) (Graamans et al., 2017). When sizing a heating, ventilation, and air conditioning system, it is important to consider 1) the increase in humidity in the air from plant transpiration; 2) the evaporative cooling effect of plant transpiration; and 3) the resulting increase in latent and decrease in sensible heat that has to be removed from the growing environment.

Resource Management

Resource cycling and environmental footprint of CEA in UA. Recent LCA studies in urban settings provide insights in the carbon footprint of several CE systems. Sanyé-Mengual et al. (2015) report that the environmental impact of a rooftop greenhouse in a city is higher at the construction stage and lower during the cultivation and transportation stages compared with a multi-span greenhouse in a rural area. Goldstein et al. (2016) concludes that the overall environmental impact of urban CEA depends on the local energy costs and production materials, and suggest that CEA for urban food production might be better-adapted to mild climates. While results from these projects cannot be extrapolated to all locations or CEA structures, they underline the importance of breaking down environmental footprint and costs by processes in the supply chain and location.

CEA is energy intensive. Some studies conclude that the establishment and infrastructure of CEA in cities can have a high environmental and economic footprint (Coley et al., 2009; de Villiers et al., 2011; Mundler and Rumpus, 2012). In addition, the energy requirement for heating in northern climates can be high. Therefore, some debate whether reducing distance from source to consumer justifies the high-energy footprint of CEA in urban settings (Goldstein et al., 2016; Sanyé-Mengual et al., 2015; Weber and Matthews, 2008). Alternative energy resources, such as geothermal, solar, wind, and hydropower, can be an improvement over traditional carbon-based energy sources. Nonetheless, local land and infrastructure design need to be considered in the overall footprint of a production system.

Soilless culture requires less water than soil-based agriculture. However, environmental gains from the overall net effect of water consumption in UA remain to be determined, considering that the irrigation efficiency (kg·m⁻³) and indirect water use (e.g., for cooling and heating) depends on the system infrastructure, irrigation design, and location, among other production factors. The multidimensional concept of water footprint considers the whole supply chain, as well as geographical and temporal parameters. Hoekstra et al. (2011) define water footprint of a product as “the total volume of fresh water that is used directly or indirectly to produce the product.” In urban CEs, the water footprint of the whole system must be considered to measure the true impact on water resources. Rainwater and

wastewater are potential water sources for the production system. However, it is unknown how capturing rainwater in urban environments may affect the entire water system (Goldstein et al., 2016; Hoekstra et al., 2011). In addition, wastewater must be treated to reduce human health risks from pathogens, pharmaceuticals, or excess salts, which can be energy-intensive. Goldstein et al. (2016) show that the total amount of water and nutrient use per crop for production in CEA can be lower compared with conventional agriculture. Nonetheless, until indirect effects on the overall water system are accounted for, the net effect on water consumption in urban setting is unknown.

Fertilizer applications in CEA are more efficient compared with field production because of targeted applications (in time and space) and the physical ease of recirculating nutrient solutions (Goldstein et al., 2016). In urban environments, human food waste could be used as fertilizer sources for plant production (Chiew et al., 2015; Spångberg et al., 2014). However, while integrating solid and liquid waste to supply nutrients for UA is technically feasible, its implementation and technology requires maximum integration to make it economically feasible (Villaruel Walker et al., 2014).

Economic Environment

Although buying local food products may not be economically cost-effective in terms of comparative advantages, consumer groups endorse UA for a number of reasons, such as to support local economies, promote environmental sustainability, and reduce food insecurity by providing access to local, fresh food in inner-city food deserts (Ikerd, 2017; Peterson et al., 2015; Scharber and Dancs, 2015; Steele, 2017). Although interest in UA is growing, few studies evaluate its cost effectiveness. For example, studies compare rooftop greenhouses with other rooftop production systems, but not to the cost of greenhouse production in rural areas (Quagraine et al., 2018; Tokunaga et al., 2015). There is a need for research that accurately assesses the return on investment and LCA of various types and sizes of CE systems for UA (Al-Kodmany, 2018).

The number of urban farms is currently low due to the many challenges faced by urban growers, such as tight regulations, zoning restrictions, limitations to water access, and high capital and operating costs (e.g., land and electricity), among others (Reisman, 2012; Steele, 2017). In addition, shade from tall buildings and skyscrapers is particularly problematic with growing plants in cities, partly explaining the increasing interest in rooftop greenhouses in urban settings. However, rooftop greenhouses have higher capital costs compared with standard ground-based greenhouses. An alternative to overcoming issues associated with shade is to produce in PFs, but this increases production costs even further because of the need for adequate environmental control (Banerjee and Adenauer, 2014). These high costs have

kept the number of urban farms small. For UA to be viable, CEs must derive profits from extending the economic benefit to the local community, or by focusing on targeted sales to markets that can pay high prices (Brumfield and Singer, 2018; Singer and Brumfield, 2017).

UA can bring value to local communities by providing opportunities for strengthening social bonds, expressing and maintaining cultural heritage, and engaging in activities promoting social and political change (Steele, 2017). A social business model offers opportunities for alternate funding sources from schools, governmental programs, or donors particularly interested in education, research, or community development. Some businesses have reduced labor costs through the assistance or use of volunteers (Reisman, 2012). In addition, commercial CE businesses with an integrated educational component offer the social benefit of providing job training and skill development for individuals. For example, CEA offers students opportunities to learn about food systems, nutrition, technology, and environmental sustainability, as well as teaching skills in customer service, leadership, marketing, and fundraising (Steele, 2017).

Consumer sociodemographic background typically drives preferences and willingness to pay for local food products. Yue and Tong (2009) show that urban consumers are more likely to buy locally grown produce compared with rural consumers. Therefore, due to the high costs associated with CEs for urban food production, some urban growers target high-value niche products (e.g., microgreens and heirloom tomatoes), charging premium prices to cover the added costs of operating in the city. To increase profitability, urban growers also target high-end restaurants and supermarkets, whose customers are prepared to pay a premium price for locally-grown produce (Sace and Natividad, 2015).

Potential Improvement of Product Quality in CEA

Increasing awareness of health and wellness has generated significant interest in the nutritional composition and health-promoting compounds of fruits and vegetables. The contribution of UA to human health can be significant because urban settings may provide easier access to fresh food (Orsini et al., 2013). Fruits and vegetables are excellent sources of selected minerals, vitamins, and antioxidants, and their impacts on human health are affected by environmental and cultural factors (Bian et al., 2015; Bjorkman et al., 2011). In contrast, harmful substances such as nitrate also can accumulate in these commodities, negatively affecting product quality (Colla et al., 2018). Environmental and cultural factors influencing mineral and phytochemical content of fruits and vegetables include light (intensity, quality, and duration), CO₂ concentration, water availability, fertilization, and pH, as well as plant species, cultivar, and plant developmental stage (Bjorkman et al.,

2011; Kopsell and Kopsell, 2008; Vicente et al., 2009). Using CEs provide opportunities for improving the quality of horticultural commodities through precise control and manipulation of these factors. For example, plant phytochemicals serve functional roles in response to environmental stress, and many of the compounds protecting plant cells also protect human cells (Demmig-Adams and Adams, 2002). Thus, environments inducing plant protection can stimulate accumulation of nutritional compounds of interest to consumers.

Although high light intensities are associated with phytochemical accumulation, maintaining high light with sole-source or supplemental lighting in CEs increases operating costs and may induce photoinhibition (Bian et al., 2015). Therefore, optimum light intensities should be considered to avoid these issues while maximizing phytochemical accumulation. Increasing light intensity across the range of 100 to 400 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ increases accumulation of lutein and β -carotene in kale (*Brassica oleracea*) (Lefsrud et al., 2006) and spinach (*Spinacea oleracea*) (Li et al., 2011) grown indoor with LEDs. Similarly, increasing light intensity with supplemental HPS lamps increases nutritional quality of vegetables grown in the greenhouse. For example, increasing light intensity increases sugar content in tomato (Dorais and Gosselin, 2002) and soluble sugars in lettuce (Fu et al., 2012). Additionally, chicory (*Cichorium endivium*), lettuce, mizuna (*Brassica rapa* var. *niposinica*), and chard (*Beta vulgaris* subsp. *vulgaris*) grown at a high light intensity contain higher total phenolic content compared with those grown at lower intensities (Colonna et al., 2016).

Manipulating light spectra also affects the quality of plant products, even at low light intensities. Although various light sources are used to grow plants in CEs, the narrow-band light output produced by LEDs offer the opportunity to manipulate spectra to control flavor, pigmentation, and other consumer-desired traits in CE-grown plants. While promising, there are inconsistent results on the effects of light spectra on crop quality, partly due to the complex interactions of multiple factors, including plant genetics and environmental conditions (e.g., temperature, light intensity). For a review on the application of light spectra to manipulate product quality of key crop species see Bian et al. (2015), Carvalho and Folta (2014), Mitchell et al., (2015), and Samuoliene et al. (2017).

High CO₂ concentration can also impact plant phytochemical and nutritional compounds. Based on a meta-analysis using published literature, Dong et al. (2018) surveyed the effect of elevated CO₂ on the nutritional quality of vegetables. The authors concluded that in general, elevated CO₂ has potential to improve the quality of vegetable crops by increasing fructose, glucose, total soluble sugar, total antioxidant capacity, total phenols, total flavonoids, ascorbic acid, and calcium (Dong et al., 2018). Water and

nutrient availability also affect the quality of plant products. Crops grown under water deficit tend to have higher nutrient content than those grown under abundant water, although this may partly be due to an increase in concentration per unit dry mass (Bjorkman et al., 2011). Reducing irrigation increases glucosinolate content and total carotenoids in cabbage (*Brassica oleracea*) and pak choi (*Brassica rapa* subsp. *chinensis*), respectively, compared with well-watered plants (Hanson et al., 2009; Radovich et al., 2005). In addition, manipulating the nutrient solution in soilless culture can help regulate secondary metabolite production in fruits and vegetables (Borgognone et al., 2016; Colla et al., 2013; Fallovo et al., 2009). Similarly, controlling the EC of the nutrient solution affects the composition and concentration of fruit phytochemicals (Colla et al., 2013; Roupael et al., 2012; Wu and Kubota, 2008).

Food Safety

Food safety risks for urban food production are triggered by microbiological, chemical, or external contaminants within the system. All of these risks can be prevented, minimized or reduced through proper training, documentation, policies, and resources. The Centers for Disease Control and Prevention reported that an estimated 48 million people (one in six U.S. residents) suffer from a food-borne illness each year (Scallan et al., 2011); 128,000 are hospitalized, and 3000 die each year from foodborne diseases. Between 1998 and 2008, 46% of all food-borne illnesses reported were associated with fruits, vegetables, and nuts (Painter et al., 2013). Contamination of fruits and vegetables produced in the United States is one of the most important issues for the food industry sector (Centers for Disease Control and Prevention, 2011). After recent high-profile outbreaks associated with fresh-cut produce (Centers for Disease Control and Prevention, 2011; Food and Drug Administration, 2018), widespread scrutiny of the U.S. fruit and vegetable supply underscores a growing concern regarding the awareness of and adherence to best practices by growers and processors.

Typically, fruits and vegetables grown in CEs are perceived to have fewer potential food safety concerns than field-grown produce due to their isolation from the soil and wild animals. However, human pathogens can still be introduced into CE production systems from various sources including water, substrates, and human contact (Olaimat and Holley, 2012; Shaw et al., 2016). In addition, because CEA often uses recirculating nutrient solutions, human pathogens can be introduced into the solutions (i.e., agricultural water or nutrients) and rapidly spread throughout the production system. Research in hydroponic systems show if *Escherichia coli* O157:H7, *E.coli* non-O157 STEC, and *Salmonella* are introduced into systems over a short period of time (48 h), these microor-

ganisms can survive and multiply (Shaw et al., 2016). Orozco et al. (2008) reported that 2.8% of tomatoes harvested from a greenhouse were contaminated with *Salmonella* and 0.7% with *E. coli*, whereas other environments and materials within the greenhouse tested positive for *Salmonella* included puddles, soil, cleaning cloths, and sponges.

Food safety-relevant contamination of produce can occur during production, harvest, processing, wholesale storage, distribution, retail, and preparation (i.e., food service or in-home). Contamination also can be caused by physical aspects (i.e., substrate, water, air, harvesting/processing equipment), animals (i.e., wild or domesticated), or human handlers (Food and Drug Administration, 2018). All growing materials (e.g., substrates, seeds, fertilizers) should be obtained from vendors who can provide evidence of following Good Manufacturing Practices, including careful monitoring of personnel, physical plant and grounds maintenance, sanitary operations, equipment and utensils, processing controls, warehouse and distribution, and holding and distribution of human food for use in animal food (Food and Drug Administration, 2017).

Pathogens of greatest concern in produce include viruses (*Hepatitis A* and *norovirus*), parasites (*Cryptosporidium*), and bacteria including *Bacillus cereus*, *Clostridium* spp., *E. coli* STEC, *Listeria monocytogenes*, *Salmonella* spp., *Yersinia enterocolitica*, *Shigella* spp., and *Campylobacter* spp. (Beuchat 2002; Harris et al., 2003; Olaimat and Holley, 2012). It is important to note that *Salmonella* spp. and *E. coli* O157:H7 are consistently associated with outbreaks of food-borne illness, and *Listeria monocytogenes* has been implicated in fewer, but particularly large, produce-related outbreaks (Olaimat and Holley, 2012). In addition to microbiological food safety concerns, chemical and heavy metal concerns must be considered for urban food production. These contaminants can be introduced in the water throughout the growing cycle and can cause harm to consumers if ingested in large amounts (United States Geological Survey, 2013).

Water is an essential component in all production systems and many foodborne outbreaks are associated with water contaminated with *E.coli* O157:H7, *Salmonella*, and *Cyclospora* (Hedberg and Osterholm, 1993). Furthermore, water sources can be categorized as low risk or high risk. While municipal water sources are considered low-risk due to daily testing by cities and municipalities for *E. coli*, surface water sources (e.g., ponds, streams) are high-risk since they are not regularly tested by a government body. Therefore, the end user is responsible for testing and results are more easily influenced by watershed activities and seasonal variation. Lopez-Galvez et al. (2014) showed that the prevalence of *Salmonella* spp. was 7.7% in irrigation water samples and 62.5% in reclaimed water sources for tomatoes grown hydroponically in greenhouses.

Social Environment

Humans and social systems are an important component of the horticultural industry (Lohr and Relf, 2000; Relf and Lohr, 2003). However in 2012, there were slightly more than three million growers in the United States, which accounted for less than 2% of the population (United States Department of Agriculture, 2014). This relatively recent disconnect from food production presents questions concerning the role of horticultural systems in human well-being, particularly involving the role of UA to provide services which promote public health and other social benefits (Brown and Jameton, 2000; Relf, 1992).

CEA is an important field with many unique facets. However, to date, the role of humans, social benefits, and communities with access to CE systems has not been thoroughly studied. Although we understand that urban food production can increase food access in cities, questions remain as to what extent CEA may alter social connections, community capitals, and education in both positive and negative ways (Eigenbrod and Gruda, 2015). By extrapolating from the available literature about urban horticultural production, we hypothesize that urban CE systems can have benefits for community food security, agricultural education, and horticultural therapy. For example, integrating CEA and UA has great potential to serve extreme-poverty areas by providing job opportunities and access to fresh food that can support a healthy diet (Bohm, 2017; Specht et al., 2014). In addition, hands-on teaching tools, facilitated through demonstrations of CE systems within schools and classrooms, allow students direct access to understand the process of food crop production. Previous research has shown that school gardens positively impact student attitude toward vegetables (Lineberger and Zajicek, 2000). Furthermore, students experiencing hands-on gardening opportunities in school exhibit more positive outlooks toward school (Waliczek et al., 2001). In addition, CEA may hold benefits to increase access to horticultural therapy. In settings such as prisons (Rice and Remy, 1994) and rehabilitation centers (Wichrowski et al., 2005), horticultural therapy facilitates both psychological and physiological healing and recovery (Lewis, 1994; Simson and Straus, 1997). By incorporating CE technologies into social institutions such as prisons, hospitals, or nursing homes, we may realize additional social benefits beyond the individual connection to plants. Furthermore, increasing rehabilitation time reduce taxpayer burden for social systems and promote overall human health and wellbeing in society.

Author's Viewpoints

Long-term success and economic viability of CEs used for food production largely will depend on future trends in consumer preferences and market demand for agricultural products produced in urban settings.

The efficiency of most urban CE systems will be improved by optimizing resources to maximize crop production and quality. For example, as LED technologies continue to increase in efficacy and decrease in cost, more CE operations will adopt the technology to reduce electrical costs and/or to improve product quality. Horticultural-grade LED fixtures are expected to be classified in two main groups: 1) relatively affordable fixtures with fixed spectra, often fixed emission rate (light output), high efficacy, and simple on/off controllers; and 2) custom-made fixtures with advanced controllers to manipulate emission rate, spectral control, and/or with sensor integration. The economic value to end-users will determine fixture applicability. Nonetheless, HPS lamps will continue to be widely used for greenhouse supplemental lighting due to their lower initial capital cost and acceptable efficacy compared with LEDs (approx. 1.7 to 1.9 vs. 2.4 to 3.0 $\mu\text{mol}\cdot\text{J}^{-1}$ for some LEDs).

Future research evaluating “dynamic” control of CO_2 enrichment will help reduce the long-term adaptation to high CO_2 concentration, which over time, diminishes the positive impacts of CO_2 enrichment in CEs. Studies focused on changing concentration levels during the daytime and/or manipulating the environment to increase CO_2 uptake during specific times of the day will be helpful in furthering our understanding about the potential benefits of using CO_2 enrichment in CEs. Further studies focusing on breeding or sink-source relationships may elucidate ways in which CO_2 enrichment can increase yield by ensuring translocation of photoassimilates to the harvestable portion of plants. Humidity control will continue to be an important challenge in CE production. However, with the continued technological development of dehumidification systems (e.g., heat pumps and desiccants), mainly driven by their use in commercial buildings, their adoption in CEs is expected to increase, especially with the worldwide expansion of PFs and potential adoption by some greenhouse facilities.

The applicability of CEA in urban settings as a solution to current challenges in our food-supply chain will be context-dependent. For example, from a food safety perspective, CEs provide more physical and environmental control compared with field-based production systems. Therefore, given the risks associated with foodborne illness outbreaks coming from large-scale outdoor farms, production in CEs may offer consumers peace of mind in their produce purchases.

Although few studies have evaluated food safety risk factors that could cause a foodborne illness in CEs, food safety should be a priority in all production systems growing edible products. Education about proper fresh produce handling from farm to fork can help prevent food safety contamination within CEA systems. Considering the wide variations in production systems, food safety programs for CEA could focus

on four main areas: substrate, water and nutrient solutions, facilities, and people. Prevention of foodborne microorganisms, chemicals, and contaminants from entering into a CE system should be the focus of training and protocols. Employees within production facilities should be trained on how to properly sanitize the facilities, tools, and equipment, along with personal health and hygiene (i.e., washing stations, restrooms, clothing, sickness and injury policies). Strict visitor policies minimizing introductions of food safety risks to the crops (e.g., hand washing, disposable booties, no-touching policies) should also be implemented.

Regarding the sustainability of CEA, local and regional limitations and resource availability must determine the net environmental and economic impacts of CEs in urban environments. The integration of water, energy, nutrients, and plant production in cities could help promote the local food movement (e.g., eliminate food deserts, fresh produce with large carbon-footprints) by delivering a neutral net carbon and water footprint. However, the footprint for crop production will depend on the crop and location, factors that are not commonly considered when assessing the environmental and economic benefits of CEA for urban food production. Consideration should be given to the fact that goods already flow in and out of cities; limiting these flows may result in negative side-effects for UA. Although scientists have advanced our understanding in CEA and many discuss the potential of integrating resources in built settings, few have investigated a truly integrated system in which waste is captured and processed in situ to supply water, nutrients, and energy to local CE operations. Engineers and horticulturists must work together to improve system performance at the establishment and production stages, and design truly integrated and efficient systems. Potential areas of work include infrastructure design and materials, energy production, storage and usage, and closing water, energy, and nutrient loops.

Specific outcomes from CEA to human wellbeing, social systems, and urban communities remain to be determined. As described above, research suggests many potential benefits from integrating CEA and UA. For example, non-commercial, small-scale CE systems may become available for display and hands-on teaching in classrooms and community centers. However, to incorporate CEA into social systems, plant scientists should be actively engaged in understanding and assessing human issues in our food chain. This allows an important unanswered gap in advancing interdisciplinary social sciences and may provide research opportunities to increase urban CEA for social sustainability and community resilience. Urban areas involve a diversity of interacting human components and the future of CEA for food production must consider these important social environments.

Conclusions

UA will continue to increase in importance as populations move into cities and the demand for local food increases. Although currently it is not uncommon for UA to take place on relatively inexpensive land in the urban core (e.g., public land or city-owned properties that are economically rented to reduce the municipal maintenance burden), CE food production for UA is expected to expand to land that is significantly more expensive than rural farming land. Therefore, one of our goals as CEA researchers will be to continue exploring how to make UA as economically, socially, and environmentally sustainable as possible. For example, we will try to recommend the most effective lighting sources available as we continue to explore ways to improve the efficiency of implementing LED lighting. We will also assist growers in managing available resources to reduce negative impacts of agriculture on the urban environment. We may do this by helping growers select irrigation systems and fertilizers that will result in less leaching and runoff. This is particularly important in an urban setting so that UA does not contribute to storm-water management issues. We also hope to identify crops that are economically viable to grow in an urban environment, considering external competition and market trends. Furthermore, our research efforts should consider all energy inputs to ensure that local products truly have a lower carbon footprint compared with crops produced outdoors and shipped to the city. Lastly, we must consider that UA creates green spaces in areas with very few plants, allowing city-dwellers to connect with how their produce is grown. This human connection to plants and food is a major benefit of UA that warrants our support so that urban producers can benefit from CEA.

Literature Cited

- Acock, E., D.A. Charkes-Edwards, D.J. Fitter, D.W. Hand, J. Ludwig, J. Warren-Wilson, and A.C. Withers. 1971. The contribution of leaves from different levels within a tomato crop to canopy net photosynthesis: An experimental examination of two canopy models. *J. Expt. Bot.* 29:815–827.
- Al-Kodmany, K. 2018. The vertical farm: A review of developments and implications for the vertical city. *Buildings* 8(24):1–36.
- Argo, W.R. and P.R. Fisher. 2002. Understanding pH management for container-grown crops. Meister Publishing, Willoughby, OH.
- Barker, J.C. 1990. Effects of day and night humidity on yield and fruit quality of glasshouse tomatoes (*Lycopersicon esculentum* Mill). *J. Hort. Sci.* 65:323–331.
- Banerjee, C. and L. Adenauer. 2014. Up, up and away! The economics of vertical farming. *J. Aging Stud.* 2:40–60.
- Benis, K., I. Turan, C. Reinhart, and P. Ferrão. 2018. Putting rooftops to use—A cost-benefit analysis of food production vs. energy generation under Mediterranean climates. *Cities* 78:166–179.
- Beuchat, L.R. 2002. Ecological factors influencing survival and growth of human pathogens on

- raw fruits and vegetables. *Microbes Infect.* 4:413–423.
- Bian, Z.H., Q.C. Yang, and W.K. Liu. 2015. Effects of light quality on the accumulation of phytochemicals in vegetables produced in controlled environments: A review. *J. Sci. Food Agr.* 95:869–877.
- Bjorkman, M., I. Kligen, A.N.E. Birch, A.M. Bones, T.J.A. Bruce, T.J. Johansen, R. Meadow, J. Molmann, R. Seljasen, L.E. Smart, and D. Stewart. 2011. Phytochemicals of *Brassicaceae* in plant protection and human health - Influences of climate, environment and agronomic practice. *Phytochemistry* 72:538–556.
- Blidariu, F. and A. Grozea. 2011. Increasing the economic efficiency and sustainability of indoor fish farming by means of aquaponics: Review. *Animal Sci. Biotech.* 44:1–8.
- Blok, C., B.E. Jackson, X. Guo, P.H.B. de Visser, and L.F.M. Marcelis. 2017. Maximum plant uptakes for water, nutrients, and oxygen are not always met by irrigation rate and distribution in water-based cultivation systems. *Front. Plant Sci.* 8:562.
- Bohm, M. 2017. Urban agriculture in and on buildings in North America: The unfulfilled potential to benefit marginalized communities. *Built Environ.* 43:343–363.
- Borgognone, D., Y. Roupael, M. Cardarelli, L. Lucini, and G. Colla. 2016. Changes in biomass, mineral composition, and quality of cardoon in response to NO₃⁻:Cl⁻ ratio and nitrate deprivation from the nutrient solution. *Front. Plant Sci.* 7:978.
- Both, A.J., J.M. Frantz, and B. Bugbee. 2017. Carbon dioxide enrichment in controlled environments, p. 82–86. In: R. Lopez and E. Runkle (eds.). *Light management in controlled environments*. Meister Media Worldwide, Willoughby, OH.
- Boulard, T., A. Baille, J. Lagier, M. Mermier, and E. Vanderschmitt. 1989. Water vapour transfer in a plastic house equipped with a dehumidification heat pump. *J. Agr. Eng. Res.* 44:191–204.
- Bradfield, E.G. and C.G. Guttridge. 1979. The dependence of calcium transport and leaf tipburn in strawberry on relative humidity and nutrient solution concentration. *Ann. Bot.* 43:363–372.
- Brown, K.H. and A.L. Jameton. 2000. Public health implications of urban agriculture. *J. Public Health Policy* 21:20–39.
- Brumfield, R.G. and C. Singer. 2018. Economics of Urban Ag: Targeting high-value, niche markets or products, and adapting a social business model can help urban greenhouses derive profit. *Produce Grower* October, 18–20.
- Bunt, A.C. 1988. Media and mixes for container-grown plants. Unwin Hyman Ltd., London, UK.
- Burnett, S.E., N.S. Mattson, and K.A. Williams. 2016. Substrates and fertilizers for organic container production of herbs, vegetables, and herbaceous ornamental plants grown in greenhouse in the United States. *Scientia Hort.* 208:111–119.
- Buzby, K.M. and L.-S. Lin. 2014. Scaling aquaponic systems: Balancing plant uptake with fish output. *Aquacult. Eng.* 63:39–44.
- Calvert, A. and G. Slack. 1975. Effects of carbon dioxide enrichment on growth, development and yield of glasshouse tomatoes. I. Responses to controlled concentrations. *J. Hort. Sci.* 50:61–71.
- Campen, J.B., G.P.A. Bot, and H.F. De Zwart. 2003. Dehumidification of greenhouses at northern latitudes. *Biosyst. Eng.* 86:487–493.
- Carvalho, S.D. and K.M. Folta. 2014. Environmentally modified organisms—expanding genetic potential with light. *Crit. Rev. Plant Sci.* 33:486–508.
- Centers for Disease Control and Prevention. 2011. Investigation of outbreak of infections caused by *Salmonella* Saintpaul. 5 Dec. 2017. <<http://www.cdc.gov/salmonella/saintpaul/jalapeno/>>.
- Chagvardieff, P., T. D’aletto, and M. André. 1994. Specific effects of irradiance and CO₂ concentration doublings on productivity and mineral content in lettuce. *Adv. Space Res.* 14:269–275.
- Chassériaux, G. and O. Gaschet. 2011. A multi-function dehumidifying heat pump for greenhouses. *Acta Hort.* 893:469–476.
- Chassériaux, G., E. Chantoiseau, C. Migeon, and P.E. Bourmet. 2014. A dehumidification strategy to avoid condensation on plants and spare energy in greenhouse. *Acta Hort.* 1037:453–460.
- Chiew, Y.L., J. Spångberg, A. Bakyb, P.A. Hansson, and H. Jönssona. 2015. Environmental impact of recycling digested food waste as a fertilizer in agriculture—A case study. *Resour. Conserv. Recycling* 95:1–14.
- Choi, J.H., G.C. Chung, and S.R. Suh. 1997. Effect of night humidity on the vegetative growth and the mineral composition of tomato and strawberry plants. *Scientia Hort.* 70:293–299.
- Clausen, A., H.M. Maersk-Moeller, J. Corfixen Soerensen, B.N. Joergensen, K.H. Kjaer, and C.O. Ottosen. 2015. Integrating commercial greenhouses in the smart grid with demand response based control of supplemental lighting. *Proc. Intl. Conf. Ind. Technol. Mgt. Sci.* 199–213.
- Cockshull, K.E. 1992. Crop environments. *Acta Hort.* 312:77–85.
- Coley, D.A., M. Howard, and M. Winter. 2009. Local food, food miles and carbon emissions: A comparison of farm shop and mass distribution approaches. *Food Policy* 34:150–155.
- Colla, G., H.J. Kim, M.C. Kyriacou, and Y. Roupael. 2018. Nitrate in fruits and vegetables. *Scientia Hort.* 237:221–238.
- Colla, G., Y. Roupael, M. Cardarelli, E. Svecova, E. Rea, and L. Lucini. 2013. Effects of saline stress on mineral composition, phenolics acids and flavonoids in leaves of artichoke and cardoon genotypes grown in floating system. *J. Sci. Food Agr.* 93:1119–1127.
- Colonna, E., Y. Roupael, G. Barbieri, and S. De Pascale. 2016. Nutritional quality of ten leafy vegetables harvested at two light intensities. *Food Chem.* 199:702–710.
- Coyle, B.D. and B. Ellison. 2017. Will consumers find vertically farmed produce “out of reach”? *Choices* 32:1–8.
- Davies, P.A. 2005. A solar cooling system for greenhouse food production in hot climates. *Sol. Energy* 79:661–668.
- Davis, B.E. 2011. Rooftop hospital gardens for physical therapy: A post-occupancy evaluation. *Health Environ. Res. Design J.* 4:14–43.
- de Villiers, D.S., H.C. Wien, J.E. Reid, and L.D. Albright. 2011. Energy use and yields in tomato production: Field, high tunnel and greenhouse compared for the northern tier of the USA (upstate New York). *Acta Hort.* 893:373–380.
- de Zwart, H.F. 2014. Energy conserving dehumidification of greenhouses. *Acta Hort.* 1037:203–210.
- Demmig-Adams, B. and W.W. Adams. 2002. Antioxidants in photosynthesis and human nutrition. *Science* 298:2149–2153.
- Dong, J., X. Li, W. Chu, and Z. Duan. 2017. High nitrate supply promotes nitrate assimilation and alleviates photosynthetic acclimation of cucumber plants under elevated CO₂. *Scientia Hort.* 218:275–283.
- Dong, J., N. Gruda, S.K. Lam, X. Li, and Z. Duan. 2018. Effects of elevated CO₂ on nutritional quality of vegetables: A review. *Front. Plant Sci.* 9:924.
- Dorais, M. and A. Gosselin. 2002. Physiological response of greenhouse vegetable crops to supplemental lighting. *Acta Hort.* 580:59–67.
- Dorais, M., C.A. Mitchell, and C. Kubota. 2017. Lighting greenhouse fruiting vegetables, p. 159–169. In: R. Lopez and E. Runkle (eds.). *Light management in controlled environments*. Meister Media Worldwide, Willoughby, OH.
- dos Santos, M.J.P.L. 2016. Smart cities and urban areas—Aquaponics as innovative urban agriculture. *Urban For. Urban Green.* 20:402–406.
- Dueck, T.A., J. Janse, B.A. Eveleens, F.L.K. Kempkes, and L.F.M. Marcelis. 2012. Growth of tomatoes under hybrid LED and HPS lighting. *Acta Hort.* 952:335–342.
- Eaves, J. and S. Eaves. 2018. Comparing profitability of greenhouse to a vertical farm in Quebec. *Can. J. Agr. Econ.* 66:43–54.
- Edwards-Jones, G., L.M. Canals, N. Hounsoume, M. Truninger, G. Koerber, B. Hounsoume, P. Cross, E.H. York, A. Hospido, K. Plassmann, I.M. Harris, R.T. Edwards, G.A.S. Day, A.D. Tomos, S.J. Cowell, and D.L. Jones. 2008. Testing the assertion that ‘local food is best’: The challenges of an evidence-based approach. *Trends Food Sci. Technol.* 19:265–274.
- Eigenbrod, C. and N. Gruda. 2015. Urban vegetable for food security in cities. A review. *Ag. Sust. Dev.* 35:483–498.
- Elumalai, S.D., A.M. Shaw, D.A. Pattillo, C.J. Currey, K.A. Rosentrater, and K. Xie. 2017. Influence of UV treatment on the food safety status of a model aquaponic system. *Water* 9:27.
- Enoch, H.Z., I. Rylski, and M. Spigelman. 1976. CO₂ enrichment of strawberry and cucumber plants grown in unheated greenhouses in Israel. *Scientia Hort.* 5:33–41.
- Falovo, C., Y. Roupael, M. Cardarelli, E. Rea, A. Battistelli, and G. Colla. 2009. Yield and quality of leafy lettuce in response to nutrient solution composition and growing season. *J. Food Agr. Environ.* 7:456–462.
- Faust, J.E. and J. Logan. 2018. Daily light integral: A research review and high-resolution maps of the United States. *HortScience* 53:1250–1257.
- Food and Drug Administration (FDA). 2017. Current good manufacturing practice, hazard analysis, and risk-based preventive controls for human food. 5 Aug. 2018. <<https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfcfr/CFRSearch.cfm?CFRPart=117>>.
- Food and Drug Administration (FDA) 2018. Guide to minimize food safety hazards of fresh-cut produce: Draft guidance for industry. 10 Dec. 2018. <<https://www.fda.gov/downloads/Food/GuidanceRegulation/GuidanceDocumentsRegulatoryInformation/UCM623718.pdf>>.
- Frantz, J.M., G. Ritchie, N.N. Cometti, J. Robinson, and B. Bugbee. 2004. Exploring the limits of crop productivity. Beyond the limits of tipburn in lettuce. *J. Amer. Soc. Hort. Sci.* 129:331–338.
- Fu, W., P. Li, Y. Wu, and J. Tang. 2012. Effects of different light intensities on anti-oxidative enzyme activity, quality and biomass in lettuce. *HortScience* 39:129–134.

- Fu, Y., L. Shao, H. Liu, H. Li, Z. Zhao, P. Ye, P. Chen, and H. Liu. 2015. Unexpected decrease in yield and antioxidants in vegetable at very high CO₂ levels. *Environ. Chem. Lett.* 13:473–479.
- Getter, K.L. and D.B. Rowe. 2006. The role of extensive green roofs in sustainable development. *HortScience* 41:1276–1285.
- Goldstein, B., M. Hauschild, J. Fernández, and M. Birkved. 2016. Testing the environmental performance of urban agriculture as a food supply in northern climates. *J. Clean. Prod.* 135:984–994.
- Gómez, C. and C.A. Mitchell. 2016. Physiological and productivity responses of high-wire tomato as affected by supplemental light source and distribution within the canopy. *J. Amer. Soc. Hort. Sci.* 141:196–208.
- Goto, E. and T. Takakura. 1992. Promotion of Ca accumulation in inner leaves by air supply for prevention of lettuce tipburn. *Trans. ASAE* 35:641–645.
- Graamans, L., A. van den Dobbelen, E. Meinen, and C. Stanghellini. 2017. Plant factories; crop transpiration and energy balance. *Agr. Syst.* 153:138–147.
- Guichard, S., C. Gary, C. Leonardi, and N. Bertin. 2005. Analysis of growth and water relations of tomato fruits in relation to air vapor pressure deficit and plant fruit load. *J. Plant Growth Regul.* 24:201–213.
- Gupta, D. 2017. Light emitting diodes for agriculture, Smart lighting. Springer, Singapore.
- Hanan, J.J. 1997. Greenhouses: Advanced technology for protected horticulture. CRC Press, Boca Raton, FL.
- Hanson, P., R.-Y. Yang, L.-C. Chang, L. Ledesma, and D. Ledesma. 2009. Contents of carotenoids, ascorbic acid, minerals and total glucosinolates in leafy brassica pakchoi (*Brassica rapa L. chinensis*) as affected by season and variety. *J. Sci. Food Agr.* 89:906–914.
- Harris, L.J., J.N. Farber, L.R. Beuchat, M.E. Parish, T.V. Suslow, E.H. Garrett, and F.F. Busta. 2003. Outbreaks associated with fresh produce: Incidence, growth and survival of pathogens in fresh and fresh-cut produce. *Compr. Rev. Food Sci. Food Saf.* 2:78–141.
- Hayden, A.L. 2006. Aeroponic and hydroponic systems for medicinal herb, rhizome, and root crops. *HortScience* 41:536–538.
- Hedberg, C.W. and M.T. Osterholm. 1993. Outbreaks of food-borne and waterborne viral gastroenteritis. *Clin. Microbiol. Rev.* 6:199–210.
- Heuvelink, E. and T. Kierkels. 2015. In greenhouses: Plant physiology in greenhouses. Horti-Text BV, Netherlands.
- Hodges, C.N., J.E. Groh, and A.W. Johnson. 1968. Controlled-environment agriculture for coastal desert areas. *Proc. Natl. Agr. Plastics Conf.* 8:58–68.
- Hoekstra, A.Y., A.K. Chapagain, M.M. Aldaya, and M.M. Mekonnen. 2011. The water footprint assessment manual: Setting a global standard. Earthscan, London, UK.
- Hussain, T., A.K. Verma, V.K. Tiwari, C. Prakash, G. Rathore, A.P. Shete, and K.K.T. Nuwansi. 2014. Optimizing koi carp, *Cyprinus carpio* var. Koi (*Linnaeus, 1758*), stocking density and nutrient recycling with spinach in an aquaponic system. *J. World Aquacult. Soc.* 45:652–661.
- Hussain, T., A.K. Verma, V.K. Tiwari, C. Prakash, G. Rathore, A.P. Shete, and N. Saharan. 2015. Effect of water flow rates on growth of *Cyprinus carpio* var. koi (*Cyprinus carpio L., 1758*) and spinach plant in aquaponic system. *Aquacult. Intl.* 23:369–384.
- Hynes, H.P. and G. Howe. 2004. Urban horticulture in the contemporary United States: Personal and community benefits. *Acta Hort.* 643:171–181.
- Ikerd, J. 2017. The urban agriculture revival. *J. Agr. Food Systems Community Dev.* 7:13–16.
- Jansen, G., N. Cila, M. Kanis, and Y. Slaats. 2016. Attitudes towards vertical farming at home: A user study. *Proc. CHI Conf. Human Factors in computing systems.* CHI2016:3091–3098.
- Jensen, M.H. 2002. Controlled environment agriculture in deserts, tropics, and temperate regions: A world review. *Acta Hort.* 578:19–25.
- Jones, C. and P. Gilbert. 2018. Determining the consequential life cycle greenhouse gas emissions of increased rooftop photovoltaic deployment. *J. Clean. Prod.* 184:211–219.
- Jones, J.B., Jr. 2014. Complete guide for growing plants hydroponically. CRC Press, Boca Raton, FL.
- Khater, E.G. and S.A. Ali. 2015. Effect of flow rate and length of gully on lettuce plants in aquaponic and hydroponic systems. *J. Aquac. Res. Dev.* 6:3.
- Klemencic, A.K. and T.G. Bulc. 2015. The use of vertical constructed wetland and ultrasound in aquaponic systems. *Environ. Sci. Pollut. Res. Intl.* 22:1420–1430.
- Kloas, W., R. Groß, D. Baganz, J. Graupner, H. Monsees, U. Schmidt, G. Staaks, J. Suhl, M. Tschirner, B. Wittstock, S. Wuertz, A. Zikova, and B. Rennert. 2015. A new concept for aquaponic systems to improve sustainability, increase productivity, and reduce environmental impacts. *Aquacult. Environ. Interact.* 7:179–192.
- Kopsell, D.A. and D.E. Kopsell. 2008. Genetic and environmental factors affecting plant lutein/zeaxanthin. *Agro Food Ind. Hi-Tech* 19:44–46.
- Körner, O. and H. Challa. 2003. Process-based humidity control regime for greenhouse crops. *Comput. Electron. Agr.* 39:173–192.
- Kotilainen, T., T.M. Robson, and R. Hernández. 2018. Light quality characterization under climate screens and shade nets for controlled-environment agriculture. *PLoS One* 13:e0199628.
- Kozai, T. 2013. Resource use efficiency of closed plant production system with artificial light: Concept, estimation and application to plant factory. *Proc. Jpn. Acad.* 89:447–461.
- Kozai, T. and G. Niu. 2016a. Plant factory as a resource-efficient closed plant production system, p. 69–90. In: T. Kozai, G. Niu, and M. Takagaki (eds.). *Plant factory: An indoor vertical farming system for efficient quality food production.* Academic Press, San Diego, CA.
- Kozai, T. and G. Niu. 2016b. Role of the plant factory with artificial lighting (PFAL) in urban areas, p. 7–33. In: T. Kozai, G. Niu, and M. Takagaki (eds.). *Plant factory: An indoor vertical farming system for efficient quality food production.* Academic Press, San Diego, CA.
- Kozai, T., W. Fang, C. Chun, Q. Yang, Y. Tong, R. Cheng, C. Kubota, and C. Lu. 2016. PFAL business and R&D in the world: Current status and perspectives, p. 35–8. In: T. Kozai, G. Niu, and M. Takagaki (eds.). *Plant factory: An indoor vertical farming system for efficient quality food production.* Academic Press, San Diego, CA.
- Kroggel, M.A. and C. Kubota. 2017. Controlled environment strategies for tipburn management in greenhouse strawberry production. *Acta Hort.* 1156:529–536.
- Kubota, C., S. Yokoi, M. Hayashi, Y. Fukuda, and S. Sase. 2006. Using ventilation-evapotranspiration-temperature-humidity (VETH) chart software for developing a strategy for evaporative cooling of semi-arid greenhouses. *Acta Hort.* 719:483–490.
- Kubota, C., M. Kroggel, A.J. Both, and M. Whalen. 2016. Does supplemental lighting make sense for my crop?—Empirical evaluations. *Acta Hort.* 1134:403–411.
- Lawson, L. 2005. City bountiful: A century of community gardening in America. University of California Press, Berkeley, CA.
- Lefsrud, M.G., D.A. Kopsell, D.E. Kopsell, and J. Curran-Celentano. 2006. Irradiance levels affect growth parameters and carotenoid pigments in kale and spinach grown in a controlled environment. *Physiol. Plant.* 127:624–631.
- Leonardi, C., A. Baile, and S. Guichard. 1999. Effects of fruit characteristics and climatic conditions on tomato transpiration in a greenhouse. *J. Hortic. Sci. Biotechnol.* 74:748–756.
- Lewis, C.A. 1994. Human health and well-being: The psychological, physiological, and sociological effects of plants on people. *Acta Hort.* 391:31–40.
- Lewis, W.M., J.H. Yopp, H.L. Schramm, and A.M. Brandenburg. 1978. Use of hydroponics to maintain quality of recirculated water in a fish culture system. *Trans. Amer. Fish. Soc.* 107:92–99.
- Li, J., S. Hikosaka, and E. Goto. 2011. Effects of light quality and photosynthetic photon flux on growth and carotenoid pigments in spinach (*Spinacia oleracea L.*). *Acta Hort.* 907:105–110.
- Lineberger, S.E. and J.M. Zajicek. 2000. School gardens: Can a hands-on teaching tool affect students' attitudes and behaviors regarding fruit and vegetables? *HortTechnology* 10:593–597.
- Lohr, V.I. and P.D. Relf. 2000. An overview of the current state of human issues in horticulture in the United States. *HortTechnology* 10:27–33.
- Lopez, R.G. and E. Runkle. 2017. Light management in controlled environments. Meister Media Worldwide, Willoughby, OH.
- Lopez-Galvez, F., A. Allende, F. Pedrero-Salcedo, J.J. Alarcon, and M.I. Gil. 2014. Safety assessment of greenhouse hydroponic tomatoes irrigated with reclaimed and surface water. *Int. J. Food Microbiol.* 191:97–102.
- Love, D.C., J.P. Fry, X. Li, E.S. Hill, L. Genello, K. Semmens, and R.E. Thompson. 2015. Commercial aquaponics production and profitability: Findings from an international survey. *Aquaculture* 435:67–74.
- Lychnos, G. and P.A. Davies. 2012. Modelling and experimental verification of a solar-powered liquid desiccant cooling system for greenhouse food production in hot climates. *Energy* 40:116–130.
- Maggini, R., C. Kiferle, and A. Pardossi. 2014. Hydroponic production of medicinal plants, p. 91–116. In: D.A.M. Pereira (ed.). *Medicinal plants: Antioxidant properties, traditional uses and conservation strategies.* Nova Science Publishers, Inc., Hauppauge, NY.
- Massa, G.D., C.A. Mitchell, J.C. Emmerich, and R.C. Morrow. 2005. Development of a reconfigurable LED plant-growth lighting system for equivalent system mass reduction in an ALS. 35th International Conference on Environmental Systems. Rome, Italy. SAE Technical Paper No. 2005-01-2955.
- Mei, L. and Y.J. Dai. 2008. A technical review on use of liquid-desiccant dehumidification for air-conditioning application. *Renew. Sustain. Energy Rev.* 12:662–689.
- Meier, S., K. Ackerman, E. Dahlgren, and X. Xu. 2013. Sustainable urban agriculture: Confirming

- viable scenarios for production. New York State Energy Research and Development Authority. 7 May 2019. <<https://www.nyserd.ny.gov/-/media/Files/Publications/Research/Environmental/Sustainable-Urban-Agriculture.pdf>>.
- Menard, C., M. Dorais, T. Hovi, and A. Gosselin. 2006. Developmental and physiological responses of tomato and cucumber to additional blue light. *Acta Hort.* 711:291–296.
- Migeon, C., A. Pierart, D. Lemesle, A. Travers, and G. Chassériaux. 2012. A dehumidifying heat pump for greenhouses. *Acta Hort.* 952:485–492.
- Mitchell, C.A. and G.W. Stutte. 2015. Sole-source lighting for controlled-environment agriculture. NASA Technical Reports. 5 Aug. 2018. <<https://ntrs.nasa.gov/archive/nasa/casi.ntrs.nasa.gov/20150009399.pdf>>.
- Mitchell, C.A., J.F. Burr, M.J. Dzakovich, C. Gómez, R. Lopez, R. Hernández, C. Kutoba, C.J. Currey, Q. Meng, E.S. Runkle, C.M. Bourguet, R.C. Murrow, and A.J. Both. 2015. Light-emitting diodes in horticulture. *Hort. Rev.* 43:1–87.
- Monsees, H., W. Kloas, and S. Wuertz. 2017. Decoupled systems on trial: Eliminating bottlenecks to improve aquaponic processes. *PLoS One* 12:e0183056.
- Morgan, K. 2009. Feeding the city: The challenge of urban food planning. *Intl. Plann. Stud.* 14:341–348.
- Mortensen, L.M. 1994. Effects of elevated CO₂ concentrations on growth and yield of eight vegetable species in a cool climate. *Scientia Hort.* 58:177–185.
- Mundler, P. and L. Rumpus. 2012. The energy efficiency of local food systems: A comparison between different modes of distribution. *Food Policy* 37:609–615.
- Nadal, A., P. Llorach-Massana, E. Cuerva, E. López-Capel, J.I. Montero, A. Josa, J. Rieradevall, and M. Royapoor. 2017. Building-integrated rooftop greenhouses: An energy and environmental assessment in the Mediterranean context. *Appl. Energy* 187:338–351.
- Nelson, J.A. and B. Bugbee. 2014. Economic analysis of greenhouse lighting: Light emitting diodes vs. High intensity discharge fixtures. *PLoS One* 9(6):e99010.
- Nelson, P.V. 2012. *Greenhouse operation and management*. 7th ed. Prentice Hall. Upper Saddle River, NJ.
- Olaimat, A.N. and R.A. Holley. 2012. Factors influencing the microbial safety of fresh produce: A review. *Food Microbiol.* 32:1–19.
- Orozco, L., L. Rico-Romero, and E.F. Escartin. 2008. Microbiological profile of greenhouses in a farm producing hydroponic tomatoes. *J. Food Prot.* 71:60–65.
- Orsini, F., R. Kahane, R. Nono-Womdim, and G. Gianquinto. 2013. Urban agriculture in the developing world: A review. *Agron. Sustain. Dev.* 33:695–720.
- Painter, J.A., R.M. Hoekstra, T. Ayers, R.V. Tauxe, C.R. Braden, F.J. Angulo, and P.M. Griffin. 2013. Attribution of foodborne illnesses, hospitalizations, and deaths to food commodities by using outbreak data, United States, 1998–2008. *Emerg. Infect. Dis.* 19:407–415.
- Palm, H.W., U. Knaus, S. Appelbaum, S. Goddek, S.M. Strauch, T. Vermeulen, M.H. Jijakli, and B. Kotzen. 2018. Towards commercial aquaponics: A review of systems, designs, scales, and nomenclature. *Aquacult. Intl.* 26:813–842.
- Pantanello, E., M. Cardarelli, E. Di Mattia, and G. Colla. 2015. Aquaponics and food safety: Effects of UV sterilization on total coliforms and lettuce production. *Acta Hort.* 1062:71–76.
- Papadopoulos, A.P., X. Luo, S. Leonhart, A. Gosselin, K. Pedneault, P. Angers, L. Gaudreau, and M. Dorais. 2001. Soilless greenhouse production of medicinal plants in North Eastern Canada. *Acta Hort.* 554:297–304.
- Park, M.H. and Y.B. Lee. 2001. Effects of CO₂ concentration, light intensity and nutrient level on growth of leaf lettuce in a plant factory. *Acta Hort.* 548:377–384.
- Paxton, A. 1994. *The food miles report: The dangers of long distance food transport*. Safe Alliance, London, UK.
- Peet, M.M. 1986. Acclimation to High CO₂ in Monoecious Cucumbers. *Plant Physiol.* 80:59–62.
- Peet, M.M. and D.H. Willits. 1984. CO₂ enrichment of greenhouse tomatoes using a closed-loop heat storage: Effects of cultivar and nitrogen. *Scientia Hort.* 24:21–32.
- Pérez-López, U., J. Miranda-Apodaca, M. Lacuesta, A. Mena-Petite, and A. Muñoz-Rueda. 2015. Growth and nutritional quality improvement in two differently pigmented lettuce cultivars grown under elevated CO₂ and/or salinity. *Scientia Hort.* 195:56–66.
- Peterson, H.H., M.R. Taylor, and Q. Baudouin. 2015. Preferences of locavores favoring community supported agriculture in the United States and France. *Ecol. Econ.* 119:64–73.
- Pinho, P., T. Hytönen, M. Rantanen, P. Elomaa, and L. Halonen. 2012. Dynamic control of supplemental lighting intensity in a greenhouse environment. *Light. Res. Technol.* 45:295–304.
- Poulet, L., G.D. Massa, R.C. Morrow, C.M. Bourget, R.M. Wheeler, and C.A. Mitchell. 2014. Significant reduction in energy for plant-growth lighting in space using targeted LED lighting and spectral manipulation. *Life Sci. Space Res.* 2:43–53.
- Potter, D.J. 2014. A review of the cultivation and processing of cannabis (*Cannabis sativa* L.) for production of prescription medicines in the UK. *Drug Test. Anal.* 6:31–38.
- Quagraine, K.K., R.M.V. Flores, H.-J. Kim, and V. McClain. 2018. Economic analysis of aquaponics and hydroponics production in the US midwest. *J. Appl. Aquacult.* 30:1–14.
- Radovich, T.J.K., M.D. Kleinhenz, and J.G. Streeter. 2005. Irrigation timing relative to head development influences yield components, sugar levels, and glucosinolate concentrations in cabbage. *J. Amer. Soc. Hort. Sci.* 130:943–949.
- Rakocy, J.E., M.P. Masser, and T.M. Losord. 2006. Recirculating aquaculture tank production systems: Aquaponics—integrating fish and plant culture. Southern Regional Aquaculture Center Publication no. 454.
- Rakocy, J.E., R.C. Shultz, D.S. Bailey, and E.S. Thoman. 2004. Aquaponic production of tilapia and basil: Comparing a batch and staggered cropping system. *Acta Hort.* 648:63–69.
- Raviv, M., J.H. Lieth and A. Bar-Tal. 2019. *Soiless culture: Theory and practice*. Elsevier, San Diego, CA.
- Reisman, A. 2012. *A greenhouse in the city: The uses and roles of community-oriented urban greenhouses*. Tufts University, Medford, MS Thesis.
- Relf, D. 1992. *The role of horticulture in human well-being and social development*. Timber Press, Portland, OR.
- Relf, P.D. and V.I. Lohr. 2003. Human issues in horticulture. *HortScience* 38:984–993.
- Rice, J.S. and L.L. Remy. 1994. Evaluating horticultural therapy: The ecological context of urban jail inmates. *J. Home Consumer Hort.* 1:203–224.
- Roosta, H.R. and M. Hamidpour. 2011. Effects of foliar application of some macro- and micro-nutrients on tomato plants in aquaponic and hydroponic systems. *Scientia Hort.* 129:396–402.
- Roosta, H.R. and Y. Mohsenian. 2015. Alleviation of alkalinity-induced Fe deficiency in eggplant (*Solanum melongena* L.) by foliar application of different Fe sources in recirculating system. *J. Plant Nutr.* 38:1768–1786.
- Rouphael, Y., M. Cardarelli, L. Lucini, E. Rea, and G. Colla. 2012. Nutrient solution concentration affects growth, mineral composition, phenolic acids and flavonoids in leaves of artichoke and cardoon. *HortScience* 47:1424–1429.
- Sace, C. and E.P. Natividad, Jr. 2015. Economic analysis of an urban vertical garden for hydroponic production of lettuce (*Lactuca sativa*). *Intl. J. Contemporary Appl. Sci.* 2:42–57.
- Samuoliënė, G., A. Brazaitytė, and V. Vaštakaitė. 2017. Light-Emitting Diodes (LEDs) for improved nutritional quality, p. 149–190. In: S.D. Gupta (ed.). *Light emitting diodes for agriculture: Smart lighting*. Springer Nature, Singapore.
- Sanjuan-Delmás, D., P. Llorach-Massana, A. Nadal, M. Ercilla-Montserrat, P. Muñoz, J.I. Montero, A. Josa, X. Gabarrell, and J. Rieradevall. 2018. Environmental assessment of an integrated rooftop greenhouse for food production in cities. *J. Clean. Prod.* 177:326–337.
- Sanyé-Mengual, E., P. Llorach-Masana, D. Sanjuan-Delmás, J. Oliver-Solà, A. Josa, J.I. Montero, and J. Rieradevall. 2014. The ICTA-ICP rooftop greenhouse lab (RTG-Lab): Closing metabolic flows (energy, water, CO₂) through integrated rooftop greenhouses, p. 693–701. In: R. Roggema and G. Keffe (eds.). *Finding spaces for productive cities*. VHL University of Applied Sciences, Velp, Netherlands.
- Sanyé-Mengual, E., J. Oliver-Solà, J.I. Montero, and J. Rieradevall. 2015. An environmental and economic life cycle assessment of rooftop greenhouse (RTG) implementation in Barcelona, Spain. Assessing new forms of urban agriculture from the greenhouse structure to the final product level. *Intl. J. Life Cycle Assess.* 20:350–366.
- Sanyé-Mengual, E., J. Martínez-Blanco, M. Finkbeiner, M. Cerdà, M. Camargo, A.R. Ometto, L.S. Velásquez, G. Villada, S. Niza, A. Pina, G. Ferreira, J. Oliver-Solà, J.I. Montero, and J. Rieradevall. 2018. Urban horticulture in retail parks: Environmental assessment of the potential implementation of rooftop greenhouses in European and South American cities. *J. Clean. Prod.* 172:3081–3091.
- Scallan, E., R.M. Hoekstra, F.J. Angula, R.V. Tauxe, M. Widdowson, S.L. Roy, J.L. Jones, and P.M. Griffin. 2011. Foodborne illness acquired in the United States—major pathogens. *Emerg. Infect. Dis.* 17:7–15.
- Scharber, H. and A. Dancs. 2015. Do locavores have a dilemma? Economic discourse and the local food critique. *Agr. Human Values* 33:121–133.
- Shamshiri, R.R., J.W. Jones, K.R. Thorp, D. Ahmad, H.C. Man, and S. Taheri. 2018. Review of optimum temperature, humidity, and vapour pressure deficit for microclimate evaluation and control in greenhouse cultivation of tomato: A review. *Intl. Agrophys.* 32:287–302.
- Shaw, A., K. Helterbran, M.R. Evans, and C. Currey. 2016. Growth of *Escherichia coli* O157:H7, non-O157 Shiga toxin-producing

- Escherichia coli*, and *Salmonella* in water and hydroponic fertilizer solutions. *J. Food Prot.* 79:2179–2183.
- Short, G., C. Yue, M. Abbey, N. Anderson, N. Phelps, P. Venturelli, and Z. Vickers. 2018. Consumer preferences for aquaponic produce: Implications from an experimental auction. *Agribusiness* 34:742–755.
- Simson, S. and M. Straus. 1997. *Horticulture as therapy: Principles and practice*. CRC Press, Boca Raton, FL.
- Singer, C. and R.G. Brumfield. 2017. An economic analysis of urban agriculture. *Intl. J. Agr. Sci.* 2:42–45.
- Sonneveld, C. and W. Voogt. 2009. *Plant nutrition of greenhouse crops*. Springer, Dordrecht, Netherlands.
- Spångberg, J., P. Tidåker, and H. Jönsson. 2014. Environmental impact of recycling nutrients in human excreta to agriculture compared with enhanced wastewater treatment. *Sci. Total Environ.* 493:209–219.
- Specht, K., R. Siebert, I. Hartmann, U.B. Freisinger, M. Sawicka, A. Werner, S. Thomaier, D. Henckel, H. Walk, and A. Dierich. 2014. Urban agriculture of the future: An overview of sustainability aspects of food production in and on buildings. *Agr. Human Values* 31:33–51.
- Steele, K. 2017. *Urban farming: An introduction to urban farming, from types and benefits to strategies and regulations*. Vitalyst Health Foundation, Phoenix, AZ.
- Sun, P., N. Mantri, H. Lou, Y. Hu, D. Sun, Y. Zhu, T. Dong, and H. Lu. 2012. Effects of elevated CO₂ and temperature on yield and fruit quality of strawberry (*Fragaria × ananassa* Duch.) at two levels of nitrogen application. *PLoS One* 7(7):e41000.
- Takagaki, M., H. Hara, and T. Kozai. 2016. Micro- and mini PFALS for improving the quality of life in urban areas, p. 91–104. In: T. Kozai, G. Niu, and M. Takagaki (ed.). *Plant factory: An indoor vertical farming system for efficient quality food production*. Academic Press, San Diego, CA.
- Tokunaga, K., C. Tamaru, H. Ako, and P.S. Leung. 2015. Economics of small-scale commercial aquaponics in Hawai'i. *J. World Aquacult. Soc.* 46:20–32.
- Tyson, R.V., D.D. Treadwell, and E.H. Simonne. 2011. Opportunities and challenges to sustainability in aquaponic systems. *HortTechnology* 21:6–13.
- United States Department of Agriculture (USDA). 2014. *Farm demographics: U.S. farmers by gender, age, race, ethnicity, and more*. 22 Feb. 2018. <https://www.agcensus.usda.gov/Publications/2012/Online_Resources/Highlights/Farm_Demographics/Highlights_Farm_Demographics.pdf>.
- United States Geological Survey (USGS). 2013. *Contaminants found in groundwater*. The USGS Water Science School. Feb. 2018. <<https://water.usgs.gov/edu/groundwater-contaminants.html>>.
- Vallièrès, M., D. De Halleux, M. Dorais, and S. Bouzid. 2014. Comparison of two cooling and dehumidifying methods for a semi-closed organic tomato greenhouse. *Acta Hort.* 1037:611–616.
- van Iersel, M.W. 2017. Optimizing LED lighting in controlled environment agriculture, p. 59–80. In: S.D. Gupta (ed.). *Light emitting diodes for agriculture: Smart lighting*. Springer Nature, Singapore.
- van Iersel, M.W. and D. Gianino. 2017. An adaptive control approach for light-emitting diode lights can reduce the energy costs of supplemental lighting in greenhouses. *HortScience* 52:72–77.
- van Iersel, M.W., G. Weaver, and M.T. Martin. 2016. A chlorophyll fluorescence-based bio-feedback system to control photosynthetic lighting in controlled environment agriculture. *J. Amer. Soc. Hort. Sci.* 141:169–176.
- Vicente, A.R., G.A. Manganaris, G.O. Sozzi, and C.H. Crisosto. 2009. Nutritional quality of fruits and vegetables, p. 57–106. In: W. Florkowski, R. Shewfelt, B. Brueckner, and S. Prussia (eds.). *Postharvest handling: A systems approach*. 2nd ed. Elsevier Inc. Academic Press, San Diego, CA.
- Villaruel Walker, R.V., M.B. Beck, J.W. Hall, R.J. Dawson, and O. Heidrich. 2014. The energy-water-food nexus: Strategic analysis of technologies for transforming the urban metabolism. *J. Environ. Mgt.* 141:104–115.
- Vitiello, D. and C. Brinkley. 2014. The hidden history of food system planning. *J. Plann. Hist.* 13:91–112.
- von Caemmerer, S. and N. Baker. 2007. The biology of transpiration. From guard cells to globe. *Plant Physiol.* 143:3.
- Waliczek, T.M., J.C. Bradley, and J.M. Zajicek. 2001. The effect of school gardens on children's interpersonal relationships and attitudes toward school. *HortTechnology* 11:466–468.
- Walters, K.J. and C.J. Currey. 2015. Hydroponic greenhouse basil production: Comparing systems and cultivars. *HortTechnology* 25:645–650.
- Wang, J., Y. Tong, Q. Yang, and M. Xin. 2016. Performance of introducing outdoor cold air for cooling a plant production system with artificial light. *Front. Plant Sci.* 7:270.
- Weaver, G.M., M. van Iersel, and J.M. Velni. 2019. A photochemistry-based method for optimizing greenhouse supplemental light intensity. *Biosyst. Eng.* 182:123–137.
- Weber, C.L. and H.S. Matthews. 2008. Food-miles and the relative climate impacts of food choices in the United States. *Environ. Sci. Technol.* 42:3508–3513.
- Wichrowski, M., J. Whiteson, F. Haas, A. Mola, and M.J. Rey. 2005. Effects of horticultural therapy on mood and heart rate in patients participating in an inpatient cardiopulmonary rehabilitation program. *J. Cardiopulm. Rehabil. Prev.* 25:270–274.
- Wongkiew, S., Z. Hu, K. Chandran, J.W. Lee, and S.K. Khanal. 2017a. Nitrogen transformations in aquaponic systems: A review. *Aquacult. Eng.* 76:9–19.
- Wongkiew, S., B.N. Popp, H.J. Kim, and S.K. Khanal. 2017b. Fate of nitrogen in floating-raft aquaponic systems using natural abundance nitrogen isotopic compositions. *Intl. Biodegradation* 125:24–32.
- Wu, M. and C. Kubota. 2008. Effects of high electrical conductivity of nutrient solution and its application timing on lycopene, chlorophyll and sugar concentrations of hydroponic tomatoes during ripening. *Scientia Hort.* 116:122–129.
- Yue, C. and C. Tong. 2009. Organic or local? Investigating consumer preference for fresh produce using a choice experiment with real economic incentive. *HortScience* 44:366–371.
- Zhang, D., Q. Du, Z. Zhang, X. Jiao, X. Song, and J. Li. 2017. Vapour pressure deficit control in relation to water transport and water productivity in greenhouse tomato production during summer. *Sci. Rep.* 7:43461.
- Zou, Y.N., Z. Hu, J. Zhang, C. Guimbaud, Q.S. Wang, and Y.K. Fang. 2016. Effect of seasonal variation on nitrogen transformations in aquaponics of northern China. *Ecol. Eng.* 94:30–36.